ELSEVIER

Journal of Organometallic Chemistry 657 (2002) 281

Journal of Organometallic Chemistry

www.elsevier.com/locate/jorganchem

Author Index of Volume 657

Ali, S., 146
Archibald, S.J., 71
Auroux, A., 98

Babonneau, F., 75
Báča, J., 59
Bakhmutov, V.I., 155
Bakhmutova, E.V., 155
Barberà, G., 217
Barton, L., 40
Bauer, C., 205
Beletskaya, I.P., 267
Beltrán, H.I., 194
Bernard, S., 91
Bernard, R., 83
Bhatti, M.H., 146
Bonnetot, B., 83, 98, 107
Borrmann, T., 205
Borrós, S., 239
Bouix, J., 91
Bould, J., 256
Bradley, J.S., 71
Bragin, V.I., 163
Bregadze, V.I., 155, 163, 171, 267
Bubnov, Y.N., 123, 146

Carboni, B., 136
Cheng, F., 71
Chistyakov, A.L., 155
Clegg, W., 256
Cornu, D., 83, 91, 98
Císařová, I., 59

Dörfler, U., 205
Dozol, J.-F., 83
Duriez, C., 107

Endo, Y., 48
Epstein, L.M., 155
Eriksson, L., 267

Farfán, N., 194
Filin, A.M., 155
Flores, M.A., 232
Framery, E., 107
Franken, A., 176, 180

Gabel, D., 205
Gervais, C., 75
Goeva, L.V., 163
Goll, S.K., 273
Greatrex, R., 256, 279
Grintselev-Knyazev, G.V., 115
Grüner, B., 59, 83

Hofmann, M., 273
Hughes, A.K., 9
Hugh MacBride, J.A., 256

Kagechika, H., 48
Kaufmann, D.E., 136
Kennedy, J.D., 20, 176, 180, 205, 256, 262, 279
Kilner, C.A., 176, 180, 205, 256, 262
Kivekäs, R., 187, 224
Kivekas, R., 232
Kivekäs, R., 247
Kudinov, A.R., 115, 171

Llop, J., 232
Lobanova, I.A., 171
Londesborough, M.G.S., 256, 262
Lyssenko, K.A., 115

Macías, R., 40
Mancilla, T., 194
Masalles, C., 239
Meshcheryakov, V.I., 171
Miele, P., 83, 91, 98, 107

Núñez, R., 224, 247

O'Dowd, C., 20
Oliva, J.M., 232
Ormsby, D.L., 256, 279

Paavola, S., 187
Pastukhov, F.V., 123
Perdigon-Melon, J.A., 98
Perekalin, D.S., 115
Petrovskii, P.V., 115, 171
Plešek, J., 59

Rais, J., 59
Rath, N.P., 40
Rosair, G.M., 217

Santillan, R., 194
Selucký, P., 59
Shubina, E.S., 155
Sillanpää, R., 224, 232, 247
Sivaev, I.B., 155, 163, 267
Sjöberg, S., 163, 267
Songkram, C., 48
Stankevich, I.V., 155
Starikova, Z.A., 115, 163, 171
Štíbr, B., 3, 279

Takaishi, K., 48
Tanatani, A., 48
Teat, S.J., 256
Teixidor, F., 187, 217, 224, 232, 239, 247
Teplitskaya, L.N., 155
Thormeier, S., 136
Thornton-Pett, M., 20, 176, 180, 205, 256, 262
Timofeev, S.V., 171
Tok, O.L., 146, 171
Toury, B., 71, 98, 107
Toutois, P., 107
Tutusaus, O., 247

Vaultier, M., 1, 107
Viñas, C., 187, 217, 224, 232, 239, 247
Vincent, H., 91
Volkov, O., 40
Votinova, N.A., 163

Wade, K., 256
Welch, A.J., 217
Wolan, A., 129
Wrackmeyer, B., 3, 146

Yamaguchi, K., 48
Yamasaki, R., 48
Yampolsky, I.V., 123

Zaidlewicz, M., 129
Zamudio-Rivera, L.S., 194

Elsevier Science B.V.
PII: S0022-328X(02)01754-0

ELSEVIER

Journal of Organometallic Chemistry 657 (2002) 283–287

Journal of Organometallic Chemistry

www.elsevier.com/locate/jorganchem

Subject Index of Volume 657

Acetamide
Synthesis and derivatization of the 2-amino-*closo*-decaborate anion $[2\text{-}B_{10}H_9NH_3]^-$, 163
Alkynes
Reactivity of some poly-1-alkynylsilicon and -tin compounds towards triallylborane—routes to novel heterocycles, 146
Allylation
Allylboration of functionalized isoquinolines, 123
Allylboration
Allylboration of functionalized isoquinolines, 123
Amine
Synthesis and derivatization of the 2-amino-*closo*-decaborate anion $[2\text{-}B_{10}H_9NH_3]^-$, 163
Arylboronic acids
Palladium-catalyzed cross-coupling reactions of arylboronic acids and 2-I-*p*-carborane, 267
Asymmetric synthesis
Chiral boronates—versatile reagents in asymmetric synthesis, 136
Azanonaborane
Azanonaborane-pyridine derivatives $[(RC_5H_4N)B_8H_{11}NHR^+]$: synthesis, structure and some molecular-orbital calculations, 205
Aziridines
Allylboration of functionalized isoquinolines, 123

Biaryls
Chiral boronates—versatile reagents in asymmetric synthesis, 136
BN fibers
Pyrolysis of poly[2,4,6-tri(methylamino)borazine] and its conversion into BN fibers, 91
Bond-critical point
Modulation of the C–C distance in disubstituted 1,2-R_2-*o*-carboranes. Crystal structure of *closo* 1,2-$(SPh)_2$-1,2-$C_2B_{10}H_{10}$, 232
Bond modulation, *closo*-carboranes
Modulation of the C–C distance in disubstituted 1,2-R_2-*o*-carboranes. Crystal structure of *closo* 1,2-$(SPh)_2$-1,2-$C_2B_{10}H_{10}$, 232
Borane cluster compounds
The use of Kitaigorodskii's *Aufbau* principle in the solid-state study of crystalline borane compounds. A preliminary account, 20
Boranes
Allylboration of functionalized isoquinolines, 123
Chemistry on a metallathiaborane cluster Part 4: reactions of 11-vertex rhodathiaboranes with bidentate phosphines and their subsequent rearrangements, 40
Borazine
Synthesis and structure of 2,4,6-tris[tris(dimethylamino)silylamino]borazine: $\{[(CH_3)_2N]_3SiNH\}_3B_3N_3H_3$, 71
Boron
Porous boron nitride supports obtained from molecular precursors. Influence of the precursor formulation and of the thermal treatment on the properties of the BN ceramic, 98
Boronates
Chiral boronates—versatile reagents in asymmetric synthesis, 136
X-ray analysis and structural characterization of 2-phenyl-6-aza-1,3-dioxa-2-borabenzocyclononenones, 194
Boron cluster compounds
Macropolyhedral boron-containing cluster chemistry: Models for intermediates en route to globular and discoidal megaloborane assemblies. Structures of $[nido\text{-}B_{10}H_{12}(nido\text{-}B_5H_8)_2]$ and $[(CH_2CH_2C_5H_4N)\text{-}arachno\text{-}B_{10}H_{10}(NC_5H_4\text{-}closo\text{-}C_2B_{10}H_{10})]$ as determined by synchrotron X-ray diffraction analysis, 256
Boron clusters
Recent developments in the chemistry of the nine-vertex monocarbaboranes, 3
Boron compounds
X-ray analysis and structural characterization of 2-phenyl-6-aza-1,3-dioxa-2-borabenzocyclononenones, 194
Boronic acids
Syntheses with organoboranes. XIII. Synthesis of ω-(4-bromophenyl)alkanoic acids and their borylation, 129
Boron nitride
Boron nitride thin fibres obtained from a new copolymer borazine–tri(methylamino)borazine precursor, 107
High resolution solid state NMR investigation of various boron nitride preceramic polymers, 75
B-phenylation
Polyhedral monocarbaborane chemistry: The C,B-*para*-diphenyl monocarbadodecaborane anion $[1,12\text{-}Ph_2\text{-}closo\text{-}1\text{-}CB_{11}H_{10}]^-$, 176
Bromination
Pd(II) bromide complexes of 1,2-bis(diphenylphosphino)-1,2-dicarba-*closo*-dodecaborane. Crystal structures of $[PdBr_2(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CH_2Cl_2$, $[PdBr_{1.133}Cl_{0.867}(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CH_2Cl_2$ and $[PdBrCl_{0.541}Me_{0.459}(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CHCl_3$, 187
B-substitution
Polyhedral monocarbaborane chemistry: The C,B-*para*-diphenyl monocarbadodecaborane anion $[1,12\text{-}Ph_2\text{-}closo\text{-}1\text{-}CB_{11}H_{10}]^-$, 176

Carboranes
Functionalized cobalt bis(dicarbollide) ions as selective extraction reagents for removal of M^{2+} and M^{3+} cations from nuclear waste, crystal and molecular structures of the $[8,8'\text{-}\mu\text{-}ClP(O)(O)_2\text{<}(1,2\text{-}C_2B_9H_{10})_2\text{-}3,3'\text{-}Co]HN(C_2H_5)_3$ and $[8,8'\text{-}\mu\text{-}Et_2NP(O)(O)_2\text{<}(1,2\text{-}C_2B_9H_{10})_2\text{-}3,3'\text{-}Co]HN(CH_3)_3$, 59
Molecular construction based on icosahedral carboranes and aromatic *N*,*N*′-dimethylurea groups. Aromatic layered molecules and a transition metal complex, 48
Recent developments in the chemistry of the nine-vertex monocarbaboranes, 3
Catalysis
Porous boron nitride supports obtained from molecular precursors. Influence of the precursor formulation and of the

Elsevier Science B.V.
PII: S0022-328X(02)01755-2

thermal treatment on the properties of the BN ceramic, 98
C,B-diphenylation
Polyhedral monocarbaborane chemistry: The C,B-*para*-diphenyl monocarbadodecaborane anion [1,12-Ph_2-*closo*-1-$CB_{11}H_{10}$]$^-$, 176
Ceramization
Pyrolysis of poly[2,4,6-tri(methylamino)borazine] and its conversion into BN fibers, 91
Chirality
Functionalized cobalt bis(dicarbollide) ions as selective extraction reagents for removal of M^{2+} and M^{3+} cations from nuclear waste, crystal and molecular structures of the [8,8′-μ-ClP(O)(O)$_2$<(1,2-$C_2B_9H_{10}$)$_2$-3,3′-Co]HN(C_2H_5)$_3$ and [8,8′-μ-Et_2NP(O)(O)$_2$<(1,2-$C_2B_9H_{10}$)$_2$-3,3′-Co](HN(CH_3)$_3$), 59
Chiral metal complexes
Functionalized cobalt bis(dicarbollide) ions as selective extraction reagents for removal of M^{2+} and M^{3+} cations from nuclear waste, crystal and molecular structures of the [8,8′-μ-ClP(O)(O)$_2$<(1,2-$C_2B_9H_{10}$)$_2$-3,3′-Co]HN(C_2H_5)$_3$ and [8,8′-μ-Et_2NP(O)(O)$_2$<(1,2-$C_2B_9H_{10}$)$_2$-3,3′-Co](HN(CH_3)$_3$), 59
Closed monocarbaborane
Polyhedral monocarbaborane chemistry: The C,B-*para*-diphenyl monocarbadodecaborane anion [1,12-Ph_2-*closo*-1-$CB_{11}H_{10}$]$^-$, 176
Close-packing
The use of Kitaigorodskii's *Aufbau* principle in the solid-state study of crystalline borane compounds. A preliminary account, 20
Closo-decaborate
Synthesis and derivatization of the 2-amino-*closo*-decaborate anion [2-$B_{10}H_9NH_3$]$^-$, 163
Closo monocarbaborane series
Polyhedral monocarbaborane chemistry: The *closo*-[PhCB$_9$H$_7$]$^-$ and *closo*-[PhCB$_{10}$H$_{10}$]$^-$ anions: the two missing species in the *closo*-[PhCB$_n$H$_n$]$^-$ sequence, 180
Cluster chemistry
Macropolyhedral boron-containing cluster chemistry: A novel triple-cluster structural motif. Isolation and characterisation of contiguous twenty-vertex [(PPh_3)$_4$ClPd$_4$B$_{16}$H$_{17}$(PPh_3)$_2$], 262
Cluster's isomerization
Recent studies on RR′S·$C_2B_9H_{11}$ charge-compensated ligands: Crystal structures of 10-(S(CH_3)$_2$)-7,8-$C_2B_9H_{11}$ and 10-(S(CH_2)$_4$)-7,8-$C_2B_9H_{11}$, 247
Cluster's thermolisis
Recent studies on RR′S·$C_2B_9H_{11}$ charge-compensated ligands: Crystal structures of 10-(S(CH_3)$_2$)-7,8-$C_2B_9H_{11}$ and 10-(S(CH_2)$_4$)-7,8-$C_2B_9H_{11}$, 247
CMPO
Synthesis of $[B_{12}H_{12}]^{2-}$ based extractants and their application for the treatment of nuclear wastes, 83
Conducting organic polymers
Cobaltabisdicarbollide anion [Co($C_2B_9H_{11}$)$_2$]$^-$ as doping agent on intelligent membranes for ion capture, 239
Conformation
Molecular construction based on icosahedral carboranes and aromatic *N*,*N*′-dimethylurea groups. Aromatic layered molecules and a transition metal complex, 48
Conjuncto heteroboranes
Macropolyhedral thiaboranes with unusual *arachno* subclusters—a computational investigation, 273
C-phenylated carbaboranes
Polyhedral monocarbaborane chemistry: The *closo*-[PhCB$_9$H$_7$]$^-$ and *closo*-[PhCB$_{10}$H$_{10}$]$^-$ anions: the two missing species in the *closo*-[PhCB$_n$H$_n$]$^-$ sequence, 180
Cross-coupling
Palladium-catalyzed cross-coupling reactions of arylboronic acids and 2-I-*p*-carborane, 267
Cross-coupling reaction
Syntheses with organoboranes. XIII. Synthesis of ω-(4-bromophenyl)alkanoic acids and their borylation, 129
Cross-link
Pyrolysis of poly[2,4,6-tri(methylamino)borazine] and its conversion into BN fibers, 91
Crystal engineering
The use of Kitaigorodskii's *Aufbau* principle in the solid-state study of crystalline borane compounds. A preliminary account, 20
Crystallization
Pyrolysis of poly[2,4,6-tri(methylamino)borazine] and its conversion into BN fibers, 91
Crystal structures
Pd(II) bromide complexes of 1,2-bis(diphenylphosphino)-1,2-dicarba-*closo*-dodecaborane. Crystal structures of [$PdBr_2$(1,2-(PPh_2)$_2$-1,2-$C_2B_{10}H_{10}$)]·CH_2Cl_2, [$PdBr_{1.133}Cl_{0.867}$(1,2-(PPh_2)$_2$-1,2-$C_2B_{10}H_{10}$)]·CH_2Cl_2 and [$PdBrCl_{0.541}Me_{0.459}$(1,2-(PPh_2)$_2$-1,2-$C_2B_{10}H_{10}$)]·$CHCl_3$, 187
C-substituted carbaboranes
Polyhedral monocarbaborane chemistry: The *closo*-[PhCB$_9$H$_7$]$^-$ and *closo*-[PhCB$_{10}$H$_{10}$]$^-$ anions: the two missing species in the *closo*-[PhCB$_n$H$_n$]$^-$ sequence, 180
[3+2] Cycloadditions
Chiral boronates—versatile reagents in asymmetric synthesis, 136
Cyclopentanes
Chiral boronates—versatile reagents in asymmetric synthesis, 136

Deboronation
Retention of the B(3)–X (X = Br, I) bond in *closo*-*o*-carborane derivatives after nucleophilic attack. The first synthesis of [3-X-7-R-7,8-*nido*-$C_2B_9H_{10}$]$^-$ (X = Br, I). Crystal structure of [$HNMe_3$][3-I-7,8-*nido*-$C_2B_9H_{11}$], 217
Decahydro-*closo*-decaborate(2−) anion
Dihydrogen bonding of decahydro-*closo*-decaborate(2−) and dodecahydro-*closo*-dodecaborate(2−) anions with proton donors: experimental and theoretical investigation, 155
Density functional theory computations
Macropolyhedral thiaboranes with unusual *arachno* subclusters—a computational investigation, 273
Derivatization
Synthesis and derivatization of the 2-amino-*closo*-decaborate anion [2-$B_{10}H_9NH_3$]$^-$, 163
Dihydrogen-bonding
The use of Kitaigorodskii's *Aufbau* principle in the solid-state study of crystalline borane compounds. A preliminary account, 20
Dimethylamino
Synthesis and structure of 2,4,6-tris[tris(dimethylamino)silylamino]borazine: {[(CH_3)$_2$N]$_3$SiNH}$_3$$B_3N_3H_3$, 71
Dodecahydro-*closo*-dodeca-borate(2−) anion
Dihydrogen bonding of decahydro-*closo*-decaborate(2−) and dodecahydro-*closo*-dodecaborate(2−) anions with proton donors: experimental and theoretical investigation, 155

Ferracarboranes
Polysubstituted derivatives of 3-(η^5-Cp)-4-SMe_2-3,1,2-$FeC_2B_9H_{10}$. X-ray structure of 3-(η^5-Cp)-4-SMe_2-7,8-(CF_3COO)$_2$-12-HgCl-3,1,2-$FeC_2B_9H_7$, 171
Fibre
Boron nitride thin fibres obtained from a new copolymer borazine–tri(methylamino)borazine precursor, 107
Fusion modes
Macropolyhedral boron-containing cluster chemistry: A novel triple-cluster structural motif. Isolation and characterisation of contiguous twenty-vertex [(PPh_3)$_4$ClPd$_4$B$_{16}$H$_{17}$(PPh_3)$_2$], 262

Heterocycles
Reactivity of some poly-1-alkynylsilicon and -tin compounds

towards triallylborane—routes to novel heterocycles, 146
Hydrododecaborate
Synthesis of $[B_{12}H_{12}]^{2-}$ based extractants and their application for the treatment of nuclear wastes, 83
Hydrogen bond
Dihydrogen bonding of decahydro-*closo*-decaborate(2−) and dodecahydro-*closo*-dodecaborate(2−) anions with proton donors: experimental and theoretical investigation, 155
Hypho cluster
Azanonaborane-pyridine derivatives $[(R'C_5H_4N)B_9H_{11}NHR']$: synthesis, structure and some molecular-orbital calculations, 205
Iodinated ligands
Retention of the B(3)–X (X = Br, I) bond in *closo*-*o*-carborane derivatives after nucleophilic attack. The first synthesis of [3-X-7-R-7,8-*nido*-$C_2B_9H_{10}]^-$ (X = Br, I). Crystal structure of [$HNMe_3$][3-I-7,8-*nido*-$C_2B_9H_{11}$], 217

Ionic liquid
Syntheses with organoboranes. XIII. Synthesis of ω-(4-bromophenyl)alkanoic acids and their borylation, 129
IR spectroscopy
Dihydrogen bonding of decahydro-*closo*-decaborate(2−) and dodecahydro-*closo*-dodecaborate(2−) anions with proton donors: experimental and theoretical investigation, 155
Isoquinolines
Allylboration of functionalized isoquinolines, 123

Kitaigorodskii's *Aufbau* principle
The use of Kitaigorodskii's *Aufbau* principle in the solid-state study of crystalline borane compounds. A preliminary account, 20

Lewis basicity of the phosphines
Phosphine–boranes incorporating the carborane cluster, 224
Liquid–liquid extraction
Functionalized cobalt bis(dicarbollide) ions as selective extraction reagents for removal of M^{2+} and M^{3+} cations from nuclear waste, crystal and molecular structures of the [8,8′-μ-$ClP(O)(O)_2$<(1,2-$C_2B_9H_{10})_2$-3,3′-Co]$HN(C_2H_5)_3$ and [8,8′-μ-$Et_2NP(O)(O)_2$<(1,2-$C_2B_9H_{10})_2$-3,3′-Co]$[HN(CH_3)_3]$, 59
Synthesis of $[B_{12}H_{12}]^{2-}$ based extractants and their application for the treatment of nuclear wastes, 83

Macropolyhedra
Macropolyhedral thiaboranes with unusual *arachno* subclusters—a computational investigation, 273
Macropolyhedral
Macropolyhedral boron-containing cluster chemistry: A novel triple-cluster structural motif. Isolation and characterisation of contiguous twenty-vertex $[(PPh_3)_4ClPd_4B_{16}H_{17}(PPh_3)_2]$, 262
Macropolyhedral boron-containing cluster chemistry: Models for intermediates en route to globular and discoidal megaloborane assemblies. Structures of [*nido*-$B_{10}H_{12}$(*nido*-$B_5H_8)_2$] and [$(CH_2CH_2C_5H_4N)$-*arachno*-$B_{10}H_{10}(NC_5H_4$-*closo*-$C_2B_{10}H_{10})$] as determined by synchrotron X-ray diffraction analysis, 256
Megaloborane
Macropolyhedral boron-containing cluster chemistry: Models for intermediates en route to globular and discoidal megaloborane assemblies. Structures of [*nido*-$B_{10}H_{12}$(*nido*-$B_5H_8)_2$] and [$(CH_2CH_2C_5H_4N)$-*arachno*-$B_{10}H_{10}(NC_5H_4$-*closo*-$C_2B_{10}H_{10})$] as determined by synchrotron X-ray diffraction analysis, 256
Mercuration
Polysubstituted derivatives of 3-(η⁵-Cp)-4-SMe_2-3,1,2-$FeC_2B_9H_{10}$. X-ray structure of 3-(η⁵-Cp)-4-SMe_2-7,8-$(CF_3COO)_2$-12-HgCl-3,1,2-$FeC_2B_9H_7$, 171
Metallaborane
Macropolyhedral boron-containing cluster chemistry: A novel triple-cluster structural motif. Isolation and characterisation of contiguous twenty-vertex $[(PPh_3)_4ClPd_4B_{16}H_{17}(PPh_3)_2]$, 262
Metallaboranes
Chemistry on a metallathiaborane cluster Part 4: reactions of 11-vertex rhodathiaboranes with bidentate phosphines and their subsequent rearrangements, 40
Functionalized cobalt bis(dicarbollide) ions as selective extraction reagents for removal of M^{2+} and M^{3+} cations from nuclear waste, crystal and molecular structures of the [8,8′-μ-$ClP(O)(O)_2$<(1,2-$C_2B_9H_{10})_2$-3,3′-Co]$HN(C_2H_5)_3$ and [8,8′-μ-$Et_2NP(O)(O)_2$<(1,2-$C_2B_9H_{10})_2$-3,3′-Co]$[HN(CH_3)_3]$, 59
Metallacarboranes
Imide- and amide-supported Group 5 and 6 metallacarboranes, 9
Synthesis and structure of rhodium complexes with monoanionic carborane ligand [9-SMe_2-7,8-$C_2B_9H_{10}]^-$, 115
MO calculations
Azanonaborane-pyridine derivatives $[(R'C_5H_4N)B_9H_{11}NHR']$: synthesis, structure and some molecular-orbital calculations, 205
Molecular design
Molecular construction based on icosahedral carboranes and aromatic *N*,*N*′-dimethylurea groups. Aromatic layered molecules and a transition metal complex, 48
Monocarbaboranes
Recent developments in the chemistry of the nine-vertex monocarbaboranes, 3
Multinuclear
Reactivity of some poly-1-alkynylsilicon and -tin compounds towards triallylborane—routes to novel heterocycles, 146

nido Carborane derivatives
Retention of the B(3)–X (X = Br, I) bond in *closo*-*o*-carborane derivatives after nucleophilic attack. The first synthesis of [3-X-7-R-7,8-*nido*-$C_2B_9H_{10}]^-$ (X = Br, I). Crystal structure of [$HNMe_3$][3-I-7,8-*nido*-$C_2B_9H_{11}$], 217
Nido-carboranylmonosulfonium
Recent studies on RR′S·$C_2B_9H_{11}$ charge-compensated ligands: Crystal structures of 10-$(S(CH_3)_2)$-7,8-$C_2B_9H_{11}$ and 10-$(S(CH_2)_4)$-7,8-$C_2B_9H_{11}$, 247
Nido-carboranylphosphines
Phosphine–boranes incorporating the carborane cluster, 224
Niobium
Imide- and amide-supported Group 5 and 6 metallacarboranes, 9
Nitride
Porous boron nitride supports obtained from molecular precursors. Influence of the precursor formulation and of the thermal treatment on the properties of the BN ceramic, 98
N ligands
Azanonaborane-pyridine derivatives $[(R'C_5H_4N)B_9H_{11}NHR']$: synthesis, structure and some molecular-orbital calculations, 205
NMR
Reactivity of some poly-1-alkynylsilicon and -tin compounds towards triallylborane—routes to novel heterocycles, 146
X-ray analysis and structural characterization of 2-phenyl-6-aza-1,3-dioxa-2-borabenzocyclononenones, 194
NMR spectroscopy
Chemistry on a metallathiaborane cluster Part 4: reactions of 11-vertex rhodathiaboranes with bidentate phosphines and their subsequent rearrangements, 40
Dihydrogen bonding of decahydro-*closo*-decaborate(2−) and dodecahydro-*closo*-dodecaborate(2−) anions with proton donors: experimental and theoretical investigation, 155
Nuclear waste
Synthesis of $[B_{12}H_{12}]^{2-}$ based extractants and their application for the treatment of nuclear wastes, 83

o-Carboranes
Pd(II) bromide complexes of 1,2-bis(diphenylphosphino)-1,2-dicarba-*closo*-dodecaborane. Crystal structures of $[PdBr_2(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CH_2Cl_2$, $[PdBr_{1.133}Cl_{0.867}(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CH_2Cl_2$ and $[PdBrCl_{0.541}Me_{0.459}(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CHCl_3$, 187
Organoboranes
Syntheses with organoboranes. XIII. Synthesis of ω-(4-bromophenyl)alkanoic acids and their borylation, 129
Organoboration
Reactivity of some poly-1-alkynylsilicon and -tin compounds towards triallylborane—routes to novel heterocycles, 146
Oxonium
Synthesis of $[B_{12}H_{12}]^{2-}$ based extractants and their application for the treatment of nuclear wastes, 83

Palladium
Palladium-catalyzed cross-coupling reactions of arylboronic acids and 2-I-*p*-carborane, 267
Palladium-boron cluster
Macropolyhedral boron-containing cluster chemistry: A novel triple-cluster structural motif. Isolation and characterisation of contiguous twenty-vertex $[(PPh_3)_4ClPd_4B_{16}H_{17}(PPh_3)_2]$, 262
Palladium(II)
Pd(II) bromide complexes of 1,2-bis(diphenylphosphino)-1,2-dicarba-*closo*-dodecaborane. Crystal structures of $[PdBr_2(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CH_2Cl_2$, $[PdBr_{1.133}Cl_{0.867}(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CH_2Cl_2$ and $[PdBrCl_{0.541}Me_{0.459}(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CHCl_3$, 187
p-Carborane
Palladium-catalyzed cross-coupling reactions of arylboronic acids and 2-I-*p*-carborane, 267
Phosphaboranes
Recent developments in the chemistry of the nine-vertex monocarbaboranes, 3
Phosphacarboranes
Recent developments in the chemistry of the nine-vertex monocarbaboranes, 3
Phosphorus
Functionalized cobalt bis(dicarbollide) ions as selective extraction reagents for removal of M^{2+} and M^{3+} cations from nuclear waste, crystal and molecular structures of the $[8,8'\text{-}\mu\text{-}ClP(O)(O)_2<(1,2\text{-}C_2B_9H_{10})_2\text{-}3,3'\text{-}Co]HN(C_2H_5)_3$ and $[8,8'\text{-}\mu\text{-}Et_2NP(O)(O)_2<(1,2\text{-}C_2B_9H_{10})_2\text{-}3,3'\text{-}Co]HN(CH_3)_3$, 59
Poly(aminoborane)
High resolution solid state NMR investigation of various boron nitride preceramic polymers, 75
Polyborazilene
High resolution solid state NMR investigation of various boron nitride preceramic polymers, 75
PolyMAB
Pyrolysis of poly[2,4,6-tri(methylamino)borazine] and its conversion into BN fibers, 91
Polymer–mineral transition
Pyrolysis of poly[2,4,6-tri(methylamino)borazine] and its conversion into BN fibers, 91
Polypyrrole
Cobaltabisdicarbollide anion $[Co(C_2B_9H_{11})_2]^-$ as doping agent on intelligent membranes for ion capture, 239
Powders
Porous boron nitride supports obtained from molecular precursors. Influence of the precursor formulation and of the thermal treatment on the properties of the BN ceramic, 98
Precursor
Boron nitride thin fibres obtained from a new copolymer borazine–tri(methylamino)borazine precursor, 107

Quantum chemistry calculation
Dihydrogen bonding of decahydro-*closo*-decaborate(2−) and dodecahydro-*closo*-dodecaborate(2−) anions with proton donors: experimental and theoretical investigation, 155

Rhodium
Chemistry on a metallathiaborane cluster Part 4: reactions of 11-vertex rhodathiaboranes with bidentate phosphines and their subsequent rearrangements, 40
Synthesis and structure of rhodium complexes with monoanionic carborane ligand $[9\text{-}SMe_2\text{-}7,8\text{-}C_2B_9H_{10}]^-$, 115

Schiff bases
Synthesis and derivatization of the 2-amino-*closo*-decaborate anion $[2\text{-}B_{10}H_9NH_3]^-$, 163
Selectivity
Cobaltabisdicarbollide anion $[Co(C_2B_9H_{11})_2]^-$ as doping agent on intelligent membranes for ion capture, 239
Siloles
Reactivity of some poly-1-alkynylsilicon and -tin compounds towards triallylborane—routes to novel heterocycles, 146
Silylamino
Synthesis and structure of 2,4,6-tris[tris(dimethylamino)silylamino]borazine: $\{[(CH_3)_2N]_3SiNH\}_3B_3N_3H_3$, 71
Solid solution
Pd(II) bromide complexes of 1,2-bis(diphenylphosphino)-1,2-dicarba-*closo*-dodecaborane. Crystal structures of $[PdBr_2(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CH_2Cl_2$, $[PdBr_{1.133}Cl_{0.867}(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CH_2Cl_2$ and $[PdBrCl_{0.541}Me_{0.459}(1,2\text{-}(PPh_2)_2\text{-}1,2\text{-}C_2B_{10}H_{10})]\cdot CHCl_3$, 187
Solid state NMR
High resolution solid state NMR investigation of various boron nitride preceramic polymers, 75
Stereoselectivity
Allylboration of functionalized isoquinolines, 123
Structural systematics
The use of Kitaigorodskii's *Aufbau* principle in the solid-state study of crystalline borane compounds. A preliminary account, 20
Structure
Synthesis and structure of 2,4,6-tris[tris(dimethylamino)silylamino]borazine: $\{[(CH_3)_2N]_3SiNH\}_3B_3N_3H_3$, 71
Support
Porous boron nitride supports obtained from molecular precursors. Influence of the precursor formulation and of the thermal treatment on the properties of the BN ceramic, 98
Supramolecular
The use of Kitaigorodskii's *Aufbau* principle in the solid-state study of crystalline borane compounds. A preliminary account, 20
Suzuki–Miayara reactions
Palladium-catalyzed cross-coupling reactions of arylboronic acids and 2-I-*p*-carborane, 267
Synchrotron X-ray diffraction analysis
Macropolyhedral boron-containing cluster chemistry: Models for intermediates en route to globular and discoidal megaloborane assemblies. Structures of [*nido*-$B_{10}H_{12}$(*nido*-$B_5H_9)_2$] and [$(CH_2CH_2C_5H_4N)$-*arachno*-$B_{10}H_{10}(NC_5H_4$-*closo*-$C_2B_{10}H_{10})$] as determined by synchrotron X-ray diffraction analysis, 256
Synthesis
Synthesis and structure of 2,4,6-tris[tris(dimethylamino)silylamino]borazine: $\{[(CH_3)_2N]_3SiNH\}_3B_3N_3H_3$, 71

Tantalum
Imide- and amide-supported Group 5 and 6 metallacarboranes, 9
Tertiary phosphonium salts
Phosphine–boranes incorporating the carborane cluster, 224
Thermolysis
Macropolyhedral boron-containing cluster chemistry: A novel

triple-cluster structural motif. Isolation and characterisation of contiguous twenty-vertex $[(PPh_3)_4ClPd_4B_{16}H_{17}(PPh_3)_2]$, 262

Thiaboranes

Chemistry on a metallathiaborane cluster Part 4: reactions of 11-vertex rhodathiaboranes with bidentate phosphines and their subsequent rearrangements, 40

Macropolyhedral thiaboranes with unusual *arachno* subclusters—a computational investigation, 273

Triple cluster fusion

Macropolyhedral boron-containing cluster chemistry: Models for intermediates en route to globular and discoidal megaloborane assemblies. Structures of $[nido\text{-}B_{10}H_{12}(nido\text{-}B_5H_8)_2]$ and $[(CH_2CH_2C_5H_4N)\text{-}arachno\text{-}B_{10}H_{10}(NC_5H_4\text{-}closo\text{-}C_2B_{10}H_{10})]$ as determined by synchrotron X-ray diffraction analysis, 256

Ureas

Molecular construction based on icosahedral carboranes and aromatic *N*,*N*′-dimethylurea groups. Aromatic layered molecules and a transition metal complex, 48

Weak intermolecular interactions

The use of Kitaigorodskii's *Aufbau* principle in the solid-state study of crystalline borane compounds. A preliminary account, 20

Weakly coordinating anions

Cobaltabisdicarbollide anion $[Co(C_2B_9H_{11})_2]^-$ as doping agent on intelligent membranes for ion capture, 239

X-ray crystallography

Functionalized cobalt bis(dicarbollide) ions as selective extraction reagents for removal of M^{2+} and M^{3+} cations from nuclear waste, crystal and molecular structures of the $[8,8'\text{-}\mu\text{-}ClP(O)(O)_2<(1,2\text{-}C_2B_9H_{10})_2\text{-}3,3'\text{-}Co]HN(C_2H_5)_3$ and $[8,8'\text{-}\mu\text{-}Et_2NP(O)(O)_2<(1,2\text{-}C_2B_9H_{10})_2\text{-}3,3'\text{-}Co](HN(CH_3)_3)$, 59

X-ray crystal structure

Azanonaborane-pyridine derivatives $[(R'C_5H_4N)B_8H_{11}NHR"]$: synthesis, structure and some molecular-orbital calculations, 205

X-ray diffraction

Chemistry on a metallathiaborane cluster Part 4: reactions of 11-vertex rhodathiaboranes with bidentate phosphines and their subsequent rearrangements, 40

Polysubstituted derivatives of $3\text{-}(\eta^5\text{-}Cp)\text{-}4\text{-}SMe_2\text{-}3,1,2\text{-}FeC_2B_9H_{10}$. X-ray structure of $3\text{-}(\eta^5\text{-}Cp)\text{-}4\text{-}SMe_2\text{-}7,8\text{-}(CF_3COO)_2\text{-}12\text{-}HgCl\text{-}3,1,2\text{-}FeC_2B_9H_7$, 171

X-ray structure

Synthesis and structure of rhodium complexes with monoanionic carborane ligand $[9\text{-}SMe_2\text{-}7,8\text{-}C_2B_9H_{10}]^-$, 115

X-ray structures

Macropolyhedral boron-containing cluster chemistry: Models for intermediates en route to globular and discoidal megaloborane assemblies. Structures of $[nido\text{-}B_{10}H_{12}(nido\text{-}B_5H_8)_2]$ and $[(CH_2CH_2C_5H_4N)\text{-}arachno\text{-}B_{10}H_{10}(NC_5H_4\text{-}closo\text{-}C_2B_{10}H_{10})]$ as determined by synchrotron X-ray diffraction analysis, 256

Polyhedral monocarbaborane chemistry: The C,B-*para*-diphenyl monocarbadodecaborane anion $[1,12\text{-}Ph_2\text{-}closo\text{-}1\text{-}CB_{11}H_{10}]^-$, 176

Polyhedral monocarbaborane chemistry: The $closo\text{-}[PhCB_7H_7]^-$ and $closo\text{-}[PhCB_{10}H_{10}]^-$ anions: the two missing species in the $closo\text{-}[PhCB_nH_n]^-$ sequence, 180

X-ray analysis and structural characterization of 2-phenyl-6-aza-1,3-dioxa-2-borabenzocyclononenones, 194

Zwitterionic clusters

Recent studies on $RR'S\cdot C_2B_9H_{11}$ charge-compensated ligands: Crystal structures of $10\text{-}(S(CH_3)_2)\text{-}7,8\text{-}C_2B_9H_{11}$ and $10\text{-}(S(CH_2)_4)\text{-}7,8\text{-}C_2B_9H_{11}$, 247